

AD-A058 042

UNIVERSITY OF SOUTHERN CALIFORNIA LOS ANGELES DEPT O--ETC F/G 7/2  
LIGAND AND SOLVENT EFFECTS ON EMISSION LIFETIMES OF CR(III) AMM--ETC(U)  
JUL 78 R T WALTERS, A W ADAMSON

N00014-76-C-0548

UNCLASSIFIED

TR-4

NL

1 OF 1  
ADA  
058042



END

DATE  
FILMED

10-78

DDC

LEVEL

OFFICE OF NAVAL RESEARCH

Contract N00014-76-C-0548

12 2

ADA 058042

9 TECHNICAL REPORT, NO. 4

6 Ligand and Solvent Effects on Emission Lifetimes of Cr(III) Amines,

by

10 R. Tom Walters and Arthur W. Adamson

Prepared for Publication

in the

Acta Chemica Scandinavica

12 29 P.

14 TR-4

DDC  
RECEIVED  
AUG 24 1978  
Jr A

Department of Chemistry  
University of Southern California  
Los Angeles, California 90007

11 July 1978

Reproduction in whole or in part is permitted for  
any purpose of the United States Government

Approved for Public Release; Distribution Unlimited.

361 555

78 08 21 027

elt

AD No. 1  
DDC FILE COPY

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 4	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) "Ligand and Solvent Effects on Emission Lifetimes of Cr(III) Amines".		5. TYPE OF REPORT & PERIOD COVERED Technical Report #4, 1978
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) R. Tom Walters and Arthur W. Adamson		8. CONTRACT OR GRANT NUMBER(s) N00014-76-C-0548
9. PERFORMING ORGANIZATION NAME AND ADDRESS University of Southern California Department of Chemistry Los Angeles, California 90007		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 051-609
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research (Code 472) Arlington, Va. 22217		12. REPORT DATE July, 1978
		13. NUMBER OF PAGES 27
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for Public Release; Distribution Unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) emission rules for emission phosphorescence chromium amines solvent effects		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Emission lifetimes and their temperature dependence are reported for a number of Cr(III) ammine complexes. A new set of rules governing emission from room temperature, fluid solution is proposed. A rationale for the rules is that thermal-type photosubstitution occurs from the $D_1$ state and that the rate controls the lifetime. FIRST THEXI DOUBLET		

Acta Chem. Scand. Ser. A. Reg. No.:

text

Received:

pages: 23

other

sheets: 4 Figs.

Correspondence and proofs should be sent to

Prof. A.W. Adamson, Department of Chemistry, University of Southern California, Los Angeles, California 90007.

Running titles: Walters and Adamson

Emission of Cr(III) Amines

---

Ligand and Solvent Effects on Emission Lifetimes of Cr(III) Amines

R. TOM WALTERS and ARTHUR W. ADAMSON

Department of Chemistry, University of Southern California, Los Angeles, California 90007.

Dedicated to Jannik Bjerrum on this 70th birthday

Lifetimes for emission from the first thexi doublet state,  $D_1^0$  ( $E_g$  in  $O_h$  symmetry) are reported for  $\text{Cr}(\text{bipy})_3^{3+}$ ,  $\text{Cr}(\text{en})_3^{3+}$ , cis- and trans- $\text{Cr}(\text{en})_2(\text{NCS})_2^+$ ,  $\text{Cr}(\text{en})(\text{NCS})_4^-$ ,  $\text{Cr}(\text{NH}_3)_6^{3+}$ ,  $\text{Cr}(\text{ND}_3)_6^{3+}$ ,  $\text{Cr}(\text{NH}_3)_5\text{Cl}^{2+}$ ,  $\text{Cr}(\text{NH}_3)_5(\text{NCS})^{2+}$ ,  $\text{Cr}(\text{NH}_3)_5(\text{H}_2\text{O})^{3+}$ , trans- $\text{Cr}(\text{NH}_3)_2(\text{NCS})_4^-$ , and  $\text{Cr}(\text{NCS})_6^{3-}$ , in water and in  $\text{D}_2\text{O}$ , acetonitrile, dimethylformamide, dimethylsulfoxide, and sulfolane solutions at 20°C. Values range from under 5 nsec to 6  $\mu\text{sec}$ . The temperature dependence of emission was determined for several systems, the apparent activation energies ranging from 7 to 12 kcal mol<sup>-1</sup>. Initial emission intensities also show temperature

78 08 21 027



dependence, with apparent activation energies,  $E_f$ , ranging from -2 to 4.5 kcal mol<sup>-1</sup>. It is suggested that emission lifetimes in room temperature fluid solution are controlled by the rate of chemical reaction from  $D_1^0$  and that  $E_f$  reflects the temperature dependence of the prompt intersystem crossing yield. A set of rules is proposed for estimating emission lifetimes of Cr(III) complexes under photochemical conditions.

ADDITIONAL		
ATIS	White Section	<input checked="" type="checkbox"/>
ROC	Diff Section	<input type="checkbox"/>
UNANNOUNCED		<input type="checkbox"/>
JUSTIFICATION		
BY		
DISPOSITION/AVAILABILITY CODES		
REL	AVAIL. ENC. OR SPECIAL	
A		

The ligand field theory of coordination compounds in general and of Cr(III) complexes in particular is, of course, a highly developed subject. Both historically and today the principal experimental parameters are the transition energies as measured by the positions of the band maxima.<sup>1,2</sup> Ligand fields are characterized in such terms, as in the spectrochemical and nephelauxetic series. It is relevant to the present paper to note that one of the approximations of ligand field theory has been the tacit ignoring of solvent effects on ligand field strengths. Although such effects are small, their existence was recognized early<sup>3</sup> and the subject has been investigated occasionally since.<sup>4</sup>

A new dimension was added to information about ligand field excited states as the photochemistry and photophysics of Werner type complexes developed to its present status.<sup>5</sup> The important excited states now appear to be those reached after thermal equilibration (thexi states) rather than the Franck-Condon state,  $Q_{FC}$ , associated with absorption band maxima. Figure 1 illustrates the distinction. Thexi states have a characteristic structure and reaction chemistry, including stereochemistry.<sup>6</sup> In the case of Cr(III) complexes, the best photochemically studied family of coordination compounds, the two lowest ligand field thexi states are a doublet,  $D_1^\circ$ , and a quartet,  $Q_1^\circ$  ( $^2E_g$  and  $^4T_{2g}$  in  $O_h$  symmetry). There has been much interest in determining the role that these states play in the photochemistry of Cr(III) complexes,<sup>7</sup> an interest greatly stimulated in recent years by the ability to observe emission from  $D_1^\circ$  under photochemical conditions.<sup>8</sup> Quenching studies<sup>9,10,11</sup> have suggested that reaction occurs via both states in the case of reinecke ion, trans- $Cr(NH_3)_2(NCS)_4^-$ ,  $(R^-)$ ,  $Cr(en)_3^{3+}$ , and trans- $Cr(en)_2(NCS)_2^+$ , and mostly via  $D_1^\circ$  in the case of  $Cr(bipy)_3^{3+}$ .<sup>12</sup>

We have been interested in the study of solvent and temperature effects on emission lifetimes as a means of understanding the nature of the rate controlling process. The results in the case of  $R^-$  ion suggested that this process might be that of direct chemical reaction from  $D_1^0$ .<sup>13</sup> The present paper extends the investigation to other Cr(III) complexes.



## EXPERIMENTAL

Materials. The various complexes were prepared by standard procedures, as described below. Chemicals generally were of reagent grade, including the solvents acetonitrile (AN), dimethylformamide (DMF), dimethylsulfoxide (DMSO), sulfolane, and pyridine.

$[\text{Cr}(\text{en})_3](\text{ClO}_4)_3$ . Tris(ethylenediamine)chromium(III) chloride  $\cdot 3\text{H}_2\text{O}$  was prepared according to a literature procedure.<sup>14</sup> The corresponding perchlorate salt was obtained by adding concentrated  $\text{HClO}_4$  dropwise to a cooled aqueous solution of the chloride salt, and recrystallized from a cooled aqueous solution. The spectrum in  $10^{-2}\text{M}$   $\text{HClO}_4$  was in excellent agreement with the literature.<sup>15</sup>  $\text{Trans}-[\text{Cr}(\text{en})_2(\text{NCS})_2]\text{NCS}$ . Trans-diisothiocyanatobis(ethylenediamine) chromium(III) thiocyanate was prepared from  $\text{Cr}(\text{en})_3(\text{NCS})_3$  according to a literature procedure.<sup>14</sup> This material was recrystallized as the perchlorate salt. The absorption spectrum was in excellent agreement with that of Bifano and Linck.<sup>16</sup>  $[\text{Cr}(\text{NH}_3)_6](\text{Cl})_3$ . Hexaamminechromium(III) chloride was prepared by a modified literature procedure.<sup>17</sup> Sublimed  $\text{CrCl}_3$  was added to condensed liquid  $\text{NH}_3$  to which had been added a small amount of Na metal and ferrous ammonium sulfate. A light red solid was collected upon evaporation of the excess liquid  $\text{NH}_3$ . Recrystallization of this material as the nitrate salt produced a yellow crystalline solid whose absorption spectrum was in excellent agreement with that reported in the literature.<sup>15</sup> The nitrate salt is not soluble in acetonitrile. However the trichloroacetic acid salt is quite soluble in that solvent and was prepared by a simple metathesis with the nitrate salt in aqueous  $\text{CCl}_3\text{COO}^-$  solution. The yellow crystalline solid was recrystallized from dilute aqueous trichloroacetic acid solution cooled to ice water temperature. Again the absorption spectrum was in excellent agreement



with that reported for  $\text{Cr}(\text{NH}_3)_6^{3+}$ .<sup>15</sup> Perdeuterohexaminechromium(III) nitrate,  $[\text{Cr}(\text{ND}_3)_6](\text{NO}_3)_3$ , was prepared by bringing a concentrated solution of the perproteo salt in  $\text{D}_2\text{O}$  to  $\text{pH} = 9$  for three minutes at  $25^\circ\text{C}$ , and then reacidifying with  $\text{D}_2\text{SO}_4$  to about  $\text{pH} = 1$ . Cooling the solution to ice water temperature produced a yellow solid material whose doublet absorption displayed the expected shift reported for the perdeutero salt.<sup>18</sup>

$[\text{Cr}(\text{en})(\text{NCS})_4]^-$  Ethylenediaminetetraisoithiocyanato chromate(III) was prepared in situ by adding ethylenediamine to a solution of  $\text{Cr}(\text{NCS})_6^{3-}$ .

The product solution displayed an absorption spectrum in excellent agreement with that reported for this complex.<sup>19</sup>  $\text{Trans}-[\text{Cr}(\text{NH}_3)_4(\text{NCS})_2]\text{NCS}$  was prepared as described by Kirk and Wong.<sup>20</sup>  $\text{Trans}-[\text{Cr}(\text{NH}_3)_4\text{Cl}_2]^+$ ,  $\text{trans}-[\text{Cr}(\text{NH}_3)_4\text{ClH}_2\text{O}]^{2+}$ , and  $\text{Cr}(\text{NH}_3)_5\text{H}_2\text{O}^{3+}$  were prepared as described by Hoppenjans, Hunt and Gregoire.<sup>21</sup> The deutero-aquo pentaamminechromium salt was prepared simply by dissolving the aquo species in  $\text{D}_2\text{O}$  and allowing exchange of the water protons for deuterons.  $[\text{Cr}(\text{NH}_3)_5\text{Cl}](\text{NO}_3)_2$ . Chloropentaamminechromium(III) chloride was purchased from Alfa Inorganics, Inc. The corresponding nitrate salt was prepared by recrystallizing the chloride salt from a fairly concentrated solution of  $\text{KNO}_3$ , and then again from a dilute solution of  $\text{KNO}_3$ . The dark red crystalline material displayed an absorption spectrum in excellent agreement with that in the literature.<sup>15</sup>

$\text{K}_3[\text{Cr}(\text{NCS})_6]$ . Potassium hexathiocyanatochromate was purchased from Alfa Inorganics, Inc., and recrystallized twice from aqueous solution.  $\text{Trans}-\text{K}[\text{Cr}(\text{NH}_3)_2(\text{NCS})_4]$ . Ammonium Reineckate was purchased from Matheson Coleman and Bell, and recrystallized as the potassium salt.<sup>22</sup>  $\text{Cis}[\text{Cr}(\text{en})_2(\text{NCS})_2]-\text{NCS}$  was prepared using a procedure analogous to that used to prepare the cis-tetraammine species.<sup>23</sup> The absorption spectrum is in excellent agreement with that reported in the literature.<sup>15</sup> Also the IR spectrum shows the expected absorptions for the cis isomer.<sup>24,25</sup>

Since emission properties were to be studied, low level impurities, if strongly emitting, could be important. The absence of any significant such contribution to emission was tested by checking that additional recrystallization of a compound had no effect on the emission behavior. In addition, our emission decay plots were first order over at least two lifetimes in all cases.

Equipment and procedures. The pulsed laser and detection equipment was that previously described,<sup>13</sup> as was the vidicon detector and multi-channel analyzer used in obtaining emission spectra. The procedures were also essentially as previously followed. Absorption spectra were obtained by means of a Beckman Acta recording spectrophotometer, and the emission yield of  $\text{Cr(en)}_3^{3+}$  was estimated from intensity data obtained with the use of an Aminco spectrofluorimeter. The emission spectrum was compared with that of a known, matching emitter, thionine.<sup>26</sup>

## RESULTS

Emission spectra. Figure 2 shows the emission spectra for  $\text{Cr}(\text{NH}_3)_6^{3+}$  and  $\text{Cr}(\text{NH}_3)_5(\text{H}_2\text{O})^{3+}$ , including the effects of deuteration. The former pair agree satisfactorily with those previously published.<sup>27</sup> The results for  $\text{Cr}(\text{NH}_3)_5(\text{H}_2\text{O})^{3+}$  are new.

Emission lifetimes. The data for 20°C are summarized in Table 1. Deuteration, either of coordinated ammonia in  $\text{Cr}(\text{NH}_3)_6^{3+}$  or of solvent make independent 15% to 30% increases in the  $D_1^\circ$  lifetime. Further, as shown in Fig. 3, the change in lifetime,  $\tau$ , for  $\text{Cr}(\text{NH}_3)_6^{3+}$  is linear in solvent  $\text{H}_2\text{O}$ - $\text{D}_2\text{O}$  composition, or essentially in composition of the solvation shell. We found the same behavior for  $\text{R}^-$  ion for acetonitrile (AN)-water mixtures.<sup>13</sup>

In general, solvent deuteration significantly increases the longer lifetime cases, but tends not to affect strongly the shorter lifetime ones. Thus  $\tau$  for trans- $\text{Cr}(\text{en})_2(\text{NCS})_2^+$  increases from 6.0 to 7.8  $\mu\text{sec}$  on solvent deuteration, while the  $\tau$ 's for cis- $\text{Cr}(\text{en})_2(\text{NCS})_2^+$ ,  $\text{Cr}(\text{NH}_3)_5(\text{NCS})_2^{2+}$ ,  $\text{R}^-$ , and trans- $\text{Cr}(\text{NH}_3)_4(\text{NCS})_2^+$  show little to no change.

Change to a non-aqueous solvent also affects the emission lifetime. The effects are not as large as those reported for  $\text{R}^-$  ion;<sup>13</sup> we were restricted in choices of solvent, however, because of the need to use polar solvents in order to have sufficient solubility. These complexes with  $\tau$ 's of 0.5  $\mu\text{sec}$  or more show solvent effects that are relatively small and variable, while those with small  $\tau$  tend to show greatly increased lifetimes in the non-aqueous media--note especially the cases of  $\text{R}^-$  ion and  $\text{Cr}(\text{NCS})_6^{3-}$ .



Temperature dependencies. Table II summarizes the results on temperature dependence. As illustrated in Fig. 4, emission lifetimes obey an Arrhenius plot within experimental error; we judge the reported apparent activation energies,  $E_p$ , to have an uncertainty of  $\pm 1$  kcal mol<sup>-1</sup> or less. Solvent effects are small, with the notable exception of Cr(bipyr)<sub>3</sub><sup>3+</sup>, for which the range is from 3.7 kcal mol<sup>-1</sup> in AN to 7.9 kcal mol<sup>-1</sup> in DMSO, including the data of Ref. 8.

The Arrhenius frequency factors,  $A$ , defined as  $A = (1/\tau)\exp(E_p/RT)$ , are included in Table II. The data fit approximately the same linear plot of  $\ln A$  vs.  $E_p$  as found to hold for  $R^-$  in various solvents.<sup>13</sup>

A second kind of temperature dependence is included in Table II. Extrapolation of the emission intensity to zero time, that is, to the inflection point of the rise of emission gave an initial reading  $V_o$ , on the oscilloscope trace. We can write  $V_o = \alpha I_o$ , where  $I_o$  is the absolute emission rate in E sec<sup>-1</sup> per E of excitation quanta absorbed and  $\alpha$  consists of various apparatus and geometry factors. For any one system, the variation of  $V_o$  with temperature (corrected for the  $\pm 10\%$  variations in exciting pulse energy as determined from the photodiode trace<sup>13</sup>) should also give the variation in  $I_o$ . That is,  $\alpha$  should not be temperature dependent. The  $V_o$  values could not be determined as precisely as the  $\tau$ 's but, as illustrated in Fig. 4, appear to fit the Arrhenius equation. The apparent activation energies,  $E_f$ , are given in Table II. Note that the values range from 4.6 kcal mol<sup>-1</sup> to -2.3 kcal mol<sup>-1</sup>; that is, in some cases  $V_o$  decreased with increasing temperature.

Emission yields,  $\phi$ , are given by  $I_o\tau$ . The yields, however, are not proportional to  $V_o\tau$  since  $\alpha$  should vary from one complex to the other because of variations in the fraction of the emission spectrum sampled by



the detector-monochromator system. Finally, although no strict correlation is expected, it is not surprising that our  $\tau$  values do vary approximately as do the reported relative emission yields.<sup>8</sup> As one comparison of interest, the emissions from trans-Cr(en)<sub>2</sub>(NCS)<sub>2</sub><sup>+</sup> and trans-Cr(NH<sub>3</sub>)<sub>2</sub>(NCS)<sub>4</sub><sup>+</sup> are reported<sup>8</sup> as "moderately strong" and "weak", respectively, while our lifetimes are 6  $\mu$ sec and 0.56  $\mu$ sec, again respectively. Parenthetically,  $\phi_p$ 's are small; thus for the relatively strong emitter, Cr(en)<sub>3</sub><sup>3+</sup>,  
 $\phi_p \approx 1 \times 10^{-4}$ .<sup>26</sup>

## DISCUSSION

Emission lifetimes of Cr(III) complexes at around room temperature presumably do not reflect  $k_r$ , the radiative rate constant. The pure radiative lifetimes of Cr(III) ammines are of the order of msec, as obtained from low temperature measurements.<sup>29</sup> It is conventional to assume that  $k_r$  is not highly environment (or temperature) dependent. The theory of spontaneous emission invokes the wave-mechanical, that is, electron,  $n$ -pole field of the molecule as interacting with the initial and final vibronic states,<sup>30</sup> and thus deals with intramolecular parameters. Alternatively, radiative transition probabilities may be estimated from absorption spectra,<sup>29</sup> and these last are nearly temperature and environment independent.<sup>3,4</sup> Also, of course, the experimental  $\phi_p$ 's are small, an indication that  $k_r \ll 1/\tau$ . Thus while in principle,  $1/\tau = (k_r + k_{nr} + k_{cr} + k_{bisc})$ , where nr denotes nonradiative relaxation to ground state, cr, chemical reaction directly from  $D_1^0$ , and bisc, back intersystem crossing to  $Q_1^0$ , it seems safe to neglect  $k_r$  as being much smaller than  $1/\tau$ .<sup>P</sup> We will also suppose that only some one of the remaining rate constants is important for our systems. In the case of  $R^-$  in various solvents, it was noted that the data obeyed a Barclay-Butler type of plot, that is, that a plot of  $\ln A$  vs.  $E_p$  was linear.<sup>13</sup> Such behavior implies that the set of data pertain to some single rate mechanism. On adding our present data a linear plot again obtains, although the data for  $R^-$  and for  $Cr(NCS)_6^{3-}$  lie somewhat above the best line for the other complexes. We will draw the same conclusion as before, namely that throughout the series it is some one of the rate constants  $k_{nr}$ ,  $k_{bisc}$ , of  $k_{cr}$  that is always largest.

The non-radiative process could be the determining one for some of our systems, but in others, quenching studies<sup>9-12,31</sup> indicate that the quantum yield for reaction from or through  $D_1^\circ$  is non-negligible, so that if some one rate process is to be dominant for the whole series, it is not the non-radiative one. For this and the reasons previously noted,<sup>13</sup> we will not consider the non-radiative process further.

The second possibility is that back intersystem crossing (see Fig. 1) is rate determining, that is, that  $D_1^\circ$  exits the excited state system via  $Q_1^\circ$ .  $E_p$  is now assigned to  $k_{bisc}$  and should be identified approximately with the  $(Q_1^\circ - D_1^\circ)$  energy difference,  $\Delta E^\circ$ . This difference cannot be obtained with any accuracy, but should at least be proportional to  $\Delta E_m$ ,<sup>32</sup> the energy corresponding to the difference between the absorption band maxima for the first ligand field band and the doublet band. Contrary to such expectation, however, there is little correlation between  $\Delta E_m$  and our  $\Delta E_p$  values. Thus for  $Cr(bipy)_3^{3+}$ ,  $Cr(en)_3^{3+}$ ,  $Cr(NH_3)_6^{3+}$ , trans- $Cr(en)_2(NCS)_2^+$ , trans- $Cr(NH_3)_2(NCS)_4^-$ , and  $Cr(NCS)_6^{3-}$  the respective  $\Delta E_m$  values are (in kcal mol<sup>-1</sup>)<sup>33</sup>: 23, 20, 18, 18, 17, and 13. The corresponding  $E_p$  values are: 7, 10, 10, 7.6, 10, and 12 (in AN).

While it has been inferred from quenching studies that back intersystem crossing is the principal mode of disappearance of  $D_1^\circ$  in the cases of  $Cr(en)_3^{3+}$  and trans- $Cr(en)_2(NCS)_2^+$ ,<sup>10,11</sup> the results can accommodate the alternative supposition that  $k_{cr}$  is the important rate constant (note also Ref. 31). Further, direct reaction from  $D_1^\circ$  has recently been proposed as dominant in the case of  $Cr(bipy)_3^{3+}$ .<sup>12</sup> Because of the lack of correlation between  $\Delta E_m$  and  $E_p$  and because  $k_{bisc}$  seems not to be important for some systems, we assume it is not important for the set of complexes, and turn to the third possibility.



The supposition that direct chemical reaction from  $D_1^0$  is the process which generally controls the lifetime of this state in room temperature fluid solution has interesting aspects. Some general implications of this mechanistic scenario were discussed previously,<sup>13</sup> and we consider here a possible explanation of the wide variations in  $\tau$  for our series of complexes. The sharpest contrast is perhaps between trans-Cr(en)<sub>2</sub>(NCS)<sub>2</sub><sup>+</sup>, with  $\tau$  = 6.0  $\mu$ sec, and trans-Cr(NH<sub>3</sub>)<sub>4</sub>(NCS)<sub>2</sub><sup>+</sup>, for which  $\tau$  is only 0.56  $\mu$ sec. Another is between trans- and cis-Cr(en)<sub>2</sub>(NCS)<sub>2</sub><sup>+</sup>, with  $\tau$  values of 6.0 and 0.52  $\mu$ sec, respectively. Note also that the series of complexes having six Cr-N bonds, Cr(bipy)<sub>3</sub><sup>3+</sup>, Cr(en)<sub>3</sub><sup>3+</sup>, Cr(NH<sub>3</sub>)<sub>6</sub><sup>3+</sup>, and Cr(NCS)<sub>6</sub><sup>3-</sup>, spans four orders of magnitude in  $\tau$ .

It appears possible to organize the data qualitatively by means of empirical rules, much as was done earlier for Cr(III) photochemistry.<sup>34</sup>

These Are:

Rule 1: For complexes with six equivalent Cr-L bands, the emission lifetime in room temperature fluid solution decreases with decreasing ligand field strength.

Rule 2: If two different kinds of ligands are coordinated, the emission lifetime will be relatively short<sup>35</sup> if that ligand which is preferentially substituted in the thermal reaction lies on the weak field axis of the complex.

That the above rules apply may be shown by means of the following examples. The order of  $\tau$ 's for the CrL<sub>6</sub> series is that of the spectrochemical series.<sup>2</sup> The weak field axis of trans-Cr(en)<sub>2</sub>(NCS)<sub>2</sub><sup>+</sup> is the SCN-Cr-NCS one, but the ground state reaction is aquation of ethylenediamine,<sup>16</sup> so rule 2 is violated, and a relatively long emission lifetime is predicted, as observed. In the case of trans-Cr(NH<sub>3</sub>)<sub>4</sub>(NCS)<sub>2</sub><sup>+</sup>, however, the thermal reaction is almost certainly one of thiocyanate aquation,<sup>36</sup>



and  $\tau$  is now small. Rule 2 likewise correctly predicts  $\tau$  to be small for cis-Cr(en)<sub>2</sub>(NCS)<sub>2</sub><sup>+</sup> and for the Cr(NH<sub>3</sub>)<sub>5</sub>X<sup>2+</sup> complexes. Cr(en)(NCS)<sub>4</sub><sup>-</sup> and trans-Cr(NH<sub>3</sub>)<sub>2</sub>(NCS)<sub>4</sub><sup>-</sup> conform; for both the weak field axis is SCN-Cr-NCS, and for both thiocyanate aquation is the preferred thermal reaction.<sup>19,37</sup>

Rule 2 carries a suggestion that emission lifetimes may tend to vary with solvent according to thermal reactivity. Some information is available on rates of solvolysis in non-aqueous solvents,<sup>38</sup> but more is needed. The case of Cr(bipy)<sub>3</sub><sup>3+</sup> may be special. The mechanism of the solvation reaction in water has complexities<sup>39</sup> and could be very solvent dependent, possibly correlating with our results.

A rationale for these rules is, of course, desirable, although not necessary to their practical usefulness. First, there is indication that direct excitation of the doublet absorption band leads preferentially to the thermal mode of reaction. This was demonstrated in the case of Cr(NH<sub>3</sub>)<sub>5</sub>(NCS)<sub>2</sub><sup>2+</sup>,<sup>40</sup> for example. We now assume this preference to be general, possibly because the D<sub>1</sub><sup>o</sup> state has or has readily accessible a vacant t<sub>2g</sub> orbital and a bimolecular substitution reaction of mechanism similar to that of the ground state process is thereby facilitated. It remains to assume that this facilitation occurs primarily for a ligand on the weak field axis and is more effective for a weak field than for a strong field ligand. If the facilitation coincides with the thermal mode of reaction, then reaction from D<sub>1</sub><sup>o</sup> is relatively fast and the lifetime of the state, relatively short.<sup>41</sup>

As with the earlier rules dealing with Cr(III) photolyses, should the present ones prove to be useful, more detailed theoretical explanations may be worth exploring. These might be along the lines developed by Zink,<sup>46</sup> Vanquickenborne,<sup>47</sup> and perhaps, especially by Wrighton<sup>48</sup> (and their co-workers).

The various deuteration effects should be commented on. As already noted, the shifts in the emission spectra presumably occur because of the vibronic nature of the transition. This aspect is presumably irrelevant to the deuteration effects on our emission lifetimes because of the unimportance of  $k_p$ . We assign the increases in  $\tau$  on deuteration to slower rates of reaction from  $D_1^\circ$ . Thermal substitution reaction rates, for example, are generally slower in  $D_2O$  than in  $H_2O$  as solvent.<sup>19,37</sup> The linear variation of  $\tau$  with degree of solvent deuteration seems consistent with this interpretation.

We turn lastly to the temperature dependencies of  $V_0$ . The initial emission intensity,  $I_0$ , to which  $V_0$  is proportional, is given by  $f_{isc}k_p$ , where  $f_{isc}$  is the efficiency of prompt intersystem crossing (see Fig. 1). Since  $k_p$  should not be very temperature dependent, our  $E_f$  values are presumably those for intersystem crossing. Present indications are that this process can be fast--occurring within short picoseconds.<sup>28</sup> We believe it to occur before complete thermal equilibration to  $Q_1^\circ$ .<sup>49</sup> A temperature dependence of  $f_{isc}$  thus indicates competition between intersystem crossing and thermal equilibration. Since the intersystem crossing event should not per se be temperature sensitive, we should interpret our  $E_f$  values in terms of changes in rates of thermal equilibration. To the extent that solvent fluidity is important in allowing the necessary bond length and bond geometry distortions to occur in the equilibration from  $Q_{FC}$  to  $Q_1^\circ$ , increasing temperature should increase the rate of thermal equilibration and thus decrease  $f_{isc}$ . We observe both positive and negative values for  $E_f$ , however, and other effects must be present. Reinecke's salt is soluble in a variety of solvents, and may therefore not be solvated in a highly specific way; for  $R^-$ , then, the negative  $E_f$  may reflect this solvent viscosity effect. The positively charged, ethylenediamine

and ammonia containing complexes, however, require polar solvents for solubility and the positive  $E_f$ 's may have to do with how a highly structured solvation shell is affected by temperature.

In the case of trans-Cr(en)<sub>2</sub>(NCS)<sub>2</sub><sup>+</sup> the lack of variation of  $V_0$  and hence presumably of  $f_{isc}$  with temperature raises the possibility that  $f_{isc}$  is near its ceiling, that is, unity. The possibility is not necessarily in contradiction to the quenching data which, in this case, indicated that about 80% (not 100%) of chemical reaction was quenched on quenching the emission from  $D_1^\circ$ .<sup>11</sup> It is quite possible that quenching encounters with  $D_1^\circ$  induce chemical reaction (in this case about 20% of the time). Thus complete emission quenching does not require complete photochemical quenching even with an  $f_{isc}$  of unity.

#### ACKNOWLEDGEMENT

These investigations were supported in part by Office of Naval Research and National Science Foundation with the University of Southern California.



Table I

## Phosphorescence Lifetimes for Cr(III) Complexes

(20°C;  $\tau$ ,  $\mu\text{sec}$ )

Complex	$\text{H}_2\text{O}$	$\text{D}_2\text{O}$	Solvent		DMSO	Sulfolane <sup>a</sup>
			AN	DMF		
$\text{Cr}(\text{bipy})_3^{3+}$ <sup>b</sup>	53 <sup>c</sup>		38			
$\text{Cr}(\text{en})_3^{3+}$	1.85	2.27	1.23	1.61	1.64	2.00
<u>trans</u> - $\text{Cr}(\text{en})_2(\text{NCS})_2^+$	6.0 <sup>d</sup>	7.8	6.3	4.75	3.6	6.5
<u>cis</u> - $\text{Cr}(\text{en})_2(\text{NCS})_2^+$	0.52	0.56	0.15	0.37	0.50	0.40
$\text{Cr}(\text{en})(\text{NCS})_4^-$	0.040 <sup>e</sup>					
$\text{Cr}(\text{NH}_3)_6^{3+}$	2.2	3.6	1.5 <sup>f</sup>	3.0	4.2	3.0
$\text{Cr}(\text{ND}_3)_6^{3+}$	3.0	3.8				
$\text{Cr}(\text{NH}_3)_5(\text{NCS})^{2+}$	0.26	0.30	g	0.51	0.92	g
<u>trans</u> - $\text{Cr}(\text{NH}_3)_4(\text{NCS})_2^+$	0.56	0.53	g	0.85	0.44	g
<u>trans</u> - $\text{Cr}(\text{NH}_3)_2(\text{NCS})_4^-$	0.005 <sup>h</sup>	0.005 <sup>h</sup>	0.12	0.080	0.090	0.25
$\text{Cr}(\text{NH}_3)_5(\text{H}_2\text{O})^{3+}$	<0.005					
$\text{Cr}(\text{NH}_3)_5(\text{D}_2\text{O})^{3+}$		<0.005				
$\text{Cr}(\text{NH}_3)_5\text{Cl}^{2+}$	<0.005					
<u>trans</u> - $\text{Cr}(\text{NH}_3)_4\text{Cl}_2^+$	<0.005				<0.005	
<u>trans</u> - $\text{Cr}(\text{NH}_3)_4(\text{H}_2\text{O})\text{Cl}^{2+}$	<0.005				<0.005	
$\text{Cr}(\text{NCS})_6^{3-}$	0.005 <sup>i</sup>		0.03	0.12	0.10	0.14

(a) glass

(b) Pyridine: 14  $\mu\text{sec}$ (c) Literature value: 51  $\mu\text{sec}$  at 22°C. <sup>12</sup>(d) Literature value: 10.5  $\mu\text{sec}$  at 15°C. <sup>11</sup>

(e) Maximum value because of possible impurities.

(f) solubilized as the  $\text{CCl}_3\text{COO}^-$  salt.

(g) Not sufficiently soluble.

(h) Extrapolated from lower temperatures. These values for  $R^-$  agree within experimental error with those previously reported. <sup>13</sup>(i) Rise in  $D_1^\circ$  absorption. <sup>28</sup>



Table II  
Apparent Activation Energies for Emission from Cr(III) Complexes<sup>a</sup>  
(in kcal mol<sup>-1</sup>)

Complex	Solvent								
	H <sub>2</sub> O			D <sub>2</sub> O			AN		
	E <sub>p</sub>	lnA	E <sub>f</sub>	E <sub>p</sub>	lnA	E <sub>f</sub>	E <sub>p</sub>	lnA	E <sub>f</sub>
Cr(bipy <sub>3</sub> ) <sup>3+</sup> <sup>b</sup>	7.0	22					3.7		
Cr(en) <sub>3</sub> <sup>3+</sup>	10	30	4.5	11.7	33	3.2			
trans-Cr(en) <sub>2</sub> (NCS) <sub>2</sub> <sup>+</sup>	7.6	25	0.0						
Cr(NH <sub>3</sub> ) <sub>6</sub> <sup>3+</sup> <sup>c</sup>	10	30	4.6	10.8	31				
Cr(ND <sub>3</sub> ) <sub>6</sub> <sup>3+</sup>	12.5	34	5.2						
trans-Cr(NH <sub>3</sub> ) <sub>2</sub> (NCS) <sub>4</sub> <sup>-</sup>	10	36	-2.3	9.0	35		10.5	34	-2.3
Cr(NCS) <sub>6</sub> <sup>3-</sup>							11.8	38	

(a) E<sub>p</sub>: from temperature dependence of the phosphorescence lifetime;

E<sub>f</sub>: from temperature dependence of V<sub>o</sub> (see text).

(b) Literature values for E<sub>p</sub> are 7.5 (H<sub>2</sub>O), 6.8 (DMF), and 7.9 (DMSO) (Ref. 8), and 9.1 (H<sub>2</sub>O) (Ref. 12). E<sub>p</sub> is 5.0 in pyridine (this work).

(c) Literature values for E<sub>p</sub> are 10.4 (H<sub>2</sub>O), 10.8 (D<sub>2</sub>O), 12.8 (DMF), 14.0 (DMSO) (Ref. 8).

## REFERENCES

1. Ballhausen, C. J., Introduction to Ligand Field Theory, Mc-Graw-Hill, New York, 1962.
2. Jørgensen, C. K., Absorption Spectra and Chemical Bonding in Complexes, Pergamon, New York, 1962.
3. Bjerrum, J., Adamson, A. W., and Bostrup, O., Acta Chem. Scand., 10, 329 (1956).
4. Adamson, A. W., J. Inorg. Nucl. Chem., 28, 1955 (1966).
5. See Adamson, A. W., and Fleischauer, P. F., Concepts of Inorganic Photochemistry, Wiley, New York, 1975.
6. Adamson, A. W., Adv. in Chem., 150, 128 (1976).
7. See Zinato, E. in Ref. 5.
8. Kane-Maguire, N. A. P., and Langford, C. H., Chem. Comm., 895 (1971).
9. Chen, S., and Porter, G., Chem. Phys. Lett., 8, 41 (1970).
10. Ballardin, R., Varani, G. V., Wasgestian, H. F., Moggi, L., and Balzani, V., J. Phys. Chem., 77, 2947 (1973).
11. Sandrini, D., Gandolfi, M. T., Moggi, L., and Balzani, V., J. Amer. Chem. Soc., 100, 1463 (1978).
12. Maestri, M., Bolletta, F., Moggi, L., Balzani, V., Henry, M. S., and Hoffman, M. Z., J. Amer. Chem. Soc., 100, 2694 (1978).
13. Gutierrez, A. R., and Adamson, A. W., J. Phys. Chem., 82, 902 (1978).
14. Rollinson, C. L. and Bailar, J. C., Jr., Inorg. Syn., 2, 198 (1946).
15. Garner, C. S. and House, D. A., Transition Metal Chemistry, 6, 69 (1970).
16. Bifano, C. and Linck, R. G., Inorg. Chem., 13, 609 (1974).
17. Oppegard, A. L. and Bailar, J. C., Jr., Inorg. Syn., 3, 153 (1950).
18. Adamson, A. W. and Dunn, T. M., J. Molec. Spectros., 18, 831 (1965).
19. Thomas, G. and Holba, V., J. Inorg. Nucl. Chem., 31, 1749 (1969).
20. Kirk, A. D. and Wong, C. F. C., Inorg. Chim. Acta, 27, 265 (1978).

21. Hoppenjans, D. W., Hunt, J. B., and Gregoire, C. R., Inorg. Chem., 7, 2506 (1968).
22. Wegner, E. and Adamson, A. W., J. Am. Chem. Soc., 88, 394 (1966).
23. Schlissenger, G. G., Inorganic Laboratory Preparations, Chemical Publishing Company, Inc., New York, 1962, p. 225. This material courtesy of Noel A. P. Kane-Maguire.
24. Baker, W. A., Jr., and Phillips, M. G., Inorg. Chem., 6, 1042 (1966).
25. McLean, J. A., Jr., Schreiner, A. F. and Laethem, A. F., J. Inorg. Nucl. Chem., 26, 1245 (1964).
26. See Haugen, G., and Hardwich, R., J. Phys. Chem., 69, 2988 (1965); Parker, C. A., and Rees, W. T., Analyst, 85, 587 (1960). These results are courtesy of R. Fukuda.
27. Long, T. V., and Penrose, D. J. B., J. Amer. Chem. Soc., 93, 632 (1971).
28. Kirk, A. D., Hoggard, T. E., Porter, G. B., Rockley, M. G., and Windsor, M. W., Chem. Phys. Lett., 37, 199 (1976).
29. See Forster, L. S., and Porter, G.B. in Ref. 5,
30. See Kuhn, H., Physical Methods of Chemistry, Vol. 1, A. Weissberger and B. Rossiter, Ed., Wiley, New York, 1972.
31. Some current results in which the rate of primary product formation from  $\text{Cr(en)}_3^{3+}$  is observed, following 530 nm laser pulse excitation, indicates that the efficiency of reaction from  $D_1^\circ$  approaches unity. Since the efficiency of reaction from  $Q_1^\circ$  is less than unity, we can rule out back intersystem crossing followed by reaction from  $Q_1^\circ$  as the important mode of disappearance of  $D_1^\circ$ .
32. See Fleischauer, P. D., Adamson, A. W., and Sartori, G., Inorg. Reaction Mechanisms, Part II, J. O. Edwards, ed., Wiley, New York, 1972.
33. Averages of the values from our spectral data and those in Ref. 8.
34. Adamson, A. W., J. Phys. Chem., 71, 798 (1967).



35. Further study may allow the term "relatively short" to be made more quantitative. For our group of complexes, "short" means under 1  $\mu$ sec.
36. Aquation of the X group is characteristic of the  $\text{trans-Cr}(\text{NH}_3)_4\text{X}_2^+$  series: See Riccieri, P., and Zinato, E., J. Amer. Chem. Soc., 97, 6071 (1975).
37. Adamson, A. W., J. Amer. Chem. Soc., 80, 3183 (1958).
38. See Palmer, D. A., and Watts, D. W., Austr. J. Chem., 21, 2895 (1968).
39. Maestri, M., Bolletta, F., Serpone, N., Moggi, L., and Balzani, V., Inorg. Chem., 15, 2048 (1976).
40. Zinato, E., Lindholm, R. D., and Adamson, A. W., J. Amer. Chem. Soc., 91, 1076 (1969). See also Martin, J. E., and Adamson, A. W., Theoretica Chimica Acta, 20, 119 (1971).
41. We plan a more thorough analysis in a subsequent publication.  $\text{Cr}(\text{CN})_6^{3-}$  may be an interesting qualified exception. Cyanide lies high in the spectrochemical series and  $\tau$  for  $\text{Cr}(\text{CN})_6^{3-}$  is quite large: in non-aqueous solvents (490  $\mu$ sec<sup>42</sup> to 540  $\mu$ sec<sup>43</sup> in DMF, 116  $\mu$ sec in DMSO,<sup>42</sup> and about 600  $\mu$ sec in AN.<sup>44</sup>) and thus consistent with rule 1. In water however,  $\tau$  is 0.1-0.2  $\mu$ sec depending on excitation wavelength.<sup>45</sup> Some special explanation clearly is required for this extremely large solvent dependence; there may be important oxygen or other impurity quenching (as with  $\text{Cr}(\text{bipy})_3^{3+}$ )<sup>12</sup>.
42. Kane-Maguire, N. A. P., Toney, C. G., Swiger, B., Adamson, A. W., and Wright, R. E., Inorg. Chimica Acta, 22, L11 (1977).
43. Wasgestian, H. F., J. Phys. Chem., 76, 1947 (1972).
44. Unpublished data plus those in Ref. 13.
45. Conti, C., and Forster, L. S., J. Amer. Chem. Soc., 99, 613 (1977).
46. Zink, J. I., J. Amer. Chem. Soc., 94, 8039 (1972).
47. Vanguickenborne, L. G., and Ceulemans, A., J. Amer. Chem. Soc., 100, 475 (1978).

48. Wrighton, M., Gray, H. B., and Hammond, G. S., Mol. Photochem., 5, 164 (1973).
49. See Ref. 13 and also Kane-Maguire, N. A. P., Phifer, J. E., and Toney, C. G., Inorg. Chem., 15, 593 (1976).

Figure 1. Energy level diagram for a Cr(III) complex.  $Q_0^0$  is the ground state quartet,  $Q_{FC}$  and  $Q_1^0$  are the Franck-Condon and the excited quartet states, respectively.  $D_1^0$  is the lowest doublet state.

Figure 2. Mass effect on  $D_1^0$  emission. A. Solid line:  $\text{Cr}(\text{NH}_3)_5(\text{H}_2\text{O})^{3+}$  in  $\text{H}_2\text{O}$ . Dashed line:  $\text{Cr}(\text{ND}_3)_5(\text{D}_2\text{O})^{3+}$  in  $\text{D}_2\text{O}$ . B. Solid line:  $\text{Cr}(\text{NH}_3)_6^{3+}$  in  $\text{H}_2\text{O}$ . Dashed line:  $\text{Cr}(\text{ND}_3)_6^{3+}$  in  $\text{D}_2\text{O}$ .

Figure 3.  $D_1^0$  emission lifetime for  $\text{Cr}(\text{NH}_3)_6^{3+}$  as a function of solvent  $\text{D}_2\text{O}$  composition at 20°C. Number of determinations shown above each point.

Figure 4. Temperature dependence of  $1/\tau$  and of  $V_0/V$  for  $\text{Cr}(\text{en})_3^{3+}$  in  $\text{H}_2\text{O}$ , open circles, and in  $\text{D}_2\text{O}$ , full circles.  $V$ , proportional to the excitation pulse energy, corrects  $V_0$  to the standard pulse.








Fig 1










Fig 4

TECHNICAL REPORT DISTRIBUTION LIST

<u>No. Copies</u>		<u>No. Copies</u>
	Office of Naval Research Arlington, Virginia 22217 Attn: Code 472	2
	Office of Naval Research Arlington, Virginia 22217 Attn: Code 102IP 1	6
	ONR Branch Office 536 S. Clark Street Chicago, Illinois 60605 Attn: Dr. Jerry Smith	1
	ONR Branch Office 715 Broadway New York, New York 10003 Attn: Scientific Dept.	1
	ONR Branch Office 1030 East Green Street Pasadena, California 91106 Attn: Dr. R. J. Marcus	1
	ONR Branch Office 760 Market Street, Rm. 447 San Francisco, California 94102 Attn: Dr. P. A. Miller	1
	ONR Branch Office 495 Summer Street Boston, Massachusetts 02210 Attn: Dr. L. H. Peebles	1
	Director, Naval Research Laboratory Washington, D.C. 20390 Attn: Code 6100	1
	The Asst. Secretary of the Navy (R&D) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1
	Commander, Naval Air Systems Command Department of the Navy Washington, D.C. 20360 Attn: Code 310C (H. Rosenwasser)	1
	Defense Documentation Center Building 5, Cameron Station Alexandria, Virginia 22314	12
	U.S. Army Research Office P.O. Box 12211 Research Triangle Park, N.C. 27709 Attn: CRD-AA-IP	1
	Naval Ocean Systems Center San Diego, California 92152 Attn: Mr. Joe McCartney	1
	Naval Weapons Center China Lake, California 93555 Attn: Head, Chemistry Division	1
	Naval Civil Engineering Laboratory Port Hueneme, California 93041 Attn: Mr. W. S. Haynes	1
	Professor O. Heinz Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
	Office of Naval Research Arlington, Virginia 22217 Attn: Dr. Richard S. Miller	1

# TECHNICAL REPORT DISTRIBUTION LIST

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. M. A. El-Sayed University of California Department of Chemistry Los Angeles, California 90024	1	Dr. G. B. Schuster University of Illinois Chemistry Department Urbana, Illinois 61801	1
Dr. M. W. Windsor Washington State University Department of Chemistry Pullman, Washington 99163	1	Dr. E. M. Eyring University of Utah Department of Chemistry Salt Lake City, Utah	1
Dr. E. R. Bernstein Colorado State University Department of Chemistry Fort Collins, Colorado 80521	1	<del>Dr. A. Adamson University of Southern California Department of Chemistry Los Angeles, California 90007</del>	<del>1</del>
Dr. C. A. Heller Naval Weapons Center Code 6059 China Lake, California 93555	1	Dr. M. S. Wrighton Massachusetts Institute of Technology Department of Chemistry Cambridge, Massachusetts 02139	1
Dr. M. H. Chisholm Princeton University Department of Chemistry Princeton, New Jersey 08540	1	Dr. M. Rauhut American Cyanamid Company Chemical Research Division Bound Brook, New Jersey 08805	1
Dr. J. R. MacDonald Naval Research Laboratory Chemistry Division Code 6110 Washington, D.C. 20375	1		